arXiv:1603.08870v1 [math.AG] 29 Mar 2016

Tropical Graph Curves

Madhusudan Manjunath*

Abstract: We study a tropical analogue of the notion of graph curves. Given a connected 3-regular graph G, we define a notion of tropical graph curve associated to G and show their existence when G is a 3-regular, three-connected planar graph. Furthermore, we realize these tropical graph curves as tropicalizations of graph curves. As an application, for smooth curves with a 3-regular, three-connected planar graph as their Berkovich skeleton, we construct canonical embeddings whose tropicalization preserves the topology of a Berkovich skeleton.

1 Introduction

Graph curves are stable curves associated to a 3-regular graph. A systematic study of their properties in terms of the underlying graph was initiated by Bayer and Eisenbud [3] in 1991. In this paper, we study tropical analogues of graph curves, that we refer to as *tropical graph curves*. We start with a brief review of graph curves before treating their tropical counterpart.

Let κ be an algebraically closed field and let G be a 3-regular, simple (no loops and no multiedges) connected graph. The graph curve X_G over κ associated to G is a totally degenerate curve with simple nodes whose dual graph is G. Hence, every irreducible component of X_G is rational. For $G = K_4$, the graph curve X_{K_4} consists of four $\mathbb{P}^1_\kappa$s with every pair intersecting transversally. The condition 3-regular ensures that X_G is well-defined up to isomorphism i.e., any choice of three distinct points as nodes on each irreducible component gives isomorphic graph curves.

The arithmetic genus of the graph curve X_G is equal to the genus of G. The graph curve X_G of arithmetic genus g carries a canonical bundle of degree $2g-2$ and rank $g-1$ [3]. It is very ample if and only if G is three-(vertex) connected [3, Proposition 2.5]. As shown in [3, Lemma 2.6], for 3-regular, simple graphs, three vertex connectivity is equivalent to three edge connectivity. We use the term *three-connected* to refer to it. For a three-connected graph G of genus g, the canonical bundle of X_G embeds into $\mathbb{P}^{g-1}_\kappa$ as an arrangement of $2g-2$ lines.

Tropical Graph Curves: For a graph G, a *geometric realization* of G is a metric graph whose underlying combinatorial graph is isomorphic to G. A *modification* of a graph G is a

*Part of this work was done while the author was at the University of California, Berkeley and was funded by a Feoder-Lynen Fellowship of the Humboldt Foundation.

metric graph G obtained by gluing (arbitrary) metric trees to a finite number of points of a geometric realization of G.

Given a connected, simple graph G of genus g, a *tropical graph curve* $\mathbb{T}_G$ associated to G is an arrangement of $2g-2$ (tropical) lines in tropical projective space $\mathbb{TP}^{g-1}$ such that $\mathbb{T}_G$ (with respect to the Euclidean topology on $\mathbb{TP}^{g-1}$, see Section 2) is homeomorphic to a modification of G.

Figure 1 shows a tropical graph curve associated to the three-dimensional cube and the eight lines corresponding to the vertices are denoted by L_{v_i}. This tropical cube curve is homeomorphic to the one-skeleton of a three-dimensional cube with trees "attached" to it at points p_1, p_2, p_3 and p_4.

The concept of tropical graph curve raises two basic questions:

1. **Existence:** Given a simple connected graph G, does G have a tropical graph curve associated to it?

2. **Realizability:** Suppose that G is a 3-regular connected graph, which tropical graph curves are realizable as the tropicalization of a canonical embedding of the graph curve X_G with respect to the trivial valuation? Three-connectivity is a necessary condition for realizability as a tropical graph curve [3, Proposition 2.5]. Canonical embeddings of graph curves whose tropicalizations yield tropical graph curves are called *weakly faithful* canonical embeddings, this terminology is justified by Section 4.

We report our results on each of the two questions. We show the existence of tropical graph curves for three-connected planar graphs.

Proposition 1.1. (**Existence Theorem**) *For every* 3*-regular three-connected planar graph* G*, there is a tropical graph curve associated to it.*

We prove Theorem 1.1 by constructing weakly faithful canonical embeddings of the corresponding graph curve. These weakly faithful canonical embeddings are derived from a planar embedding of the graph. For instance, the following arrangement of lines gives a weakly faithful tropicalization of the (one-skeleton of the) three-dimensional cube. Consider the planar embedding of the cube $\mathcal{C}$ with its facets (and vertices) labelled as shown in Figure 2. Let x_{F_j} be the variable corresponding to the face F_j and identify $\mathbb{P}^4_\kappa$ with $\mathrm{Proj}(\kappa[x_{F_1},\ldots,x_{F_5}])$. The equations for the eight lines corresponding to eight vertices are as follows:

$$\begin{gathered}
L_{v_1} = \langle x_{F_1} + x_{F_2} + x_{F_3}, x_{F_4}, x_{F_5}\rangle \\
L_{v_2} = \langle x_{F_1} + x_{F_4} + x_{F_3}, x_{F_2}, x_{F_5}\rangle \\
L_{v_3} = \langle x_{F_1} + x_{F_4} + x_{F_5}, x_{F_2}, x_{F_3}\rangle \\
L_{v_4} = \langle x_{F_1} + x_{F_2} + x_{F_5}, x_{F_3}, x_{F_4}\rangle \\
L_{v_5} = \langle x_{F_1}, x_{F_3}, x_{F_4}\rangle,\ L_{v_6} = \langle x_{F_1}, x_{F_4}, x_{F_5}\rangle \\
L_{v_7} = \langle x_{F_1}, x_{F_2}, x_{F_5}\rangle,\ L_{v_8} = \langle x_{F_1}, x_{F_2}, x_{F_3}\rangle
\end{gathered}$$



Figure 1: Tropical Cube Curve



Figure 2: A Planar Embedding of the Cube

Consider the tropicalization of this line arrangement. When seen as a subset of the algebraic torus it is not weakly faithful. In particular, the lines L_{v_5}, L_{v_6}, L_{v_7} and L_{v_8} do not intersect the algebraic torus. On the other hand, this line arrangement can be tropicalized in tropical projective space $\mathbb{TP}^4$ (this justifies the appearance of tropical projective space in the definition of tropical graph curves) to yield the tropical line arrangement shown in Figure 1. Hence, this canonical embedding is weakly faithful.

The property of being weakly faithful is highly sensitive to the choice of a canonical embedding, any linear change of co-ordinates in $\mathbb{P}^4$ yields another canonical embedding of the graph curve $X_{\mathcal{C}}$. Each of these canonical embeddings has dual graph $\mathcal{C}$ but, roughly speaking the tropicalization of "most" of these will not be weakly faithful. More precisely, by [12] there is an associated generic tropical curve and this is not typically weakly faithful.

We generalize this arrangement of lines to arbitrary 3-regular three-connected planar graphs and show the following.

Theorem 1.2. (**Realizability Theorem**) *Let G be a 3-regular, three-connected planar graph (and by Steinitz' theorem, a one-skeleton of a three dimensional polytope is earnest), there exists a canonical embedding of X_G that has a weakly faithful tropicalization.*

In fact, we construct a specific canonical embedding called the *schön embedding* (see Section 3) that has a weakly faithful tropicalization. A key ingredient in the proof of Theorem 1.1 and Theorem 1.2 is via a tropical analogue of the notion of ΔY transformation (a transformation used to prove Steinitz theorem), see Figures 9, 10 for a glimpse.

1.1 Tropical Basis for Weakly Faithful Canonical Embeddings

We describe a tropical basis for the schön embedding. It has the following simple description in terms of the Stanley-Reisner ideal of the dual complex of the polytope associated to G.

Let M be the dual complex of the three-dimensional polytope P associated to G (the existence of P is guaranteed by Steinitz theorem, see Theorem 3.8 for its statement) and let F be the set of facets of P (facets in P are in bijection with the vertices of M). The schön embedding is the Stanley-Reisner variety of M intersected with the hyperplane $\sum_{f\in F} x_f = 0$. This hyperplane section is reduced. Hence, a minimal generating set $\mathcal{G}$ of the vanishing radical ideal I_{sch} of the schön embedding is the set of monomial minimal generators of the Stanley-Reisner ideal of M with the variable x_e corresponding to the outerface face replaced by $-\sum_{f\in F, f\neq e} x_f$. Furthermore, this set is a tropical basis.

Proposition 1.3. *The set $\mathcal{G}$ forms a tropical basis for defining ideal I_{sch} of the schön embedding.*

Example 1.4. *Let $\mathcal{C}$ be the one-skeleton of the three-dimensional cube. The minimal generating set of the vanishing ideal of $X_{\mathcal{C}}$ produced by Theorem 3.1 is the following:*

$$\langle x_{F_2}x_{F_4}, x_{F_3}x_{F_5}, x_{F_1}^2 + x_{F_1}x_{F_2} + x_{F_1}x_{F_3} + x_{F_1}x_{F_4} + x_{F_1}x_{F_5}\rangle$$

1.2 Graph Curves and Tropical Curves

This work was motivated by the following potential connections between graph curves and tropical curves.

1. **Graph Curves and Tropical Curves:** The work of Bayer and Eisenbud on graph curves predates much of the literature in tropical geometry. Both tropical and graph curves are "combinatorial objects" associated to a smooth algebraic curve. From this viewpoint, it is natural to question how they relate to each other. Theorem 1.8 establishes a concrete connection between graph curves and tropical curves.

2. **Faithful Tropicalization:** Another motivation for our work comes from the concept of faithful tropicalization developed by Baker, Payne and Rabinoff [2].

   Let C be a smooth proper curve over a non-archimedean field $\mathbb{K}$ i.e., an algebraically closed field with a non-trivial non-archimedean valuation and complete with respect to it. Any tropicalization of C as well as its Berkovich analytification C^{an} have natural metrics on them ([10], [4]) and there is a continuous map from the Berkovich analytification to any tropicalization. A tropicalization of a smooth algebraic curve is called *faithful* if there exists a skeleton of its Berkovich analytification (see [2, Section 5.15.2] for a precise definition) for which this map is homeomorphic and isometric onto its image. A tropicalization is called *weakly faithful* if there exists a skeleton of the Berkovich analytification for which this map is homeomorphic onto its image.

   Theorem 1.5. (Baker, Payne and Rabinoff [2]) *Every smooth proper curve over $\mathbb{K}$ has a faithful (and hence, a weakly faithful) tropicalization.*

   The proof of Theorem 1.5 use methods from non-archimedean analysis and shows that an embedding of the curve into a split torus of sufficiently high dimension yields a faithful tropicalization. From a constructive point of view, explicitly constructing embeddings of the curve that gives faithful tropicalizations is of interest. This leads to the following problem:

   Problem 1.6. *Compute a faithful/ weakly faithful tropicalization when the curve is presented in terms of its defining equations.*

   A closely related problem is the following:

   Problem 1.7. *Explicitly describe the structure of defining equations whose tropicalization is faithful/ weakly faithful.*

   Another related problem is the following realizability question:

   Question 1. *Which graphs occur as skeleta of faithful/ weakly faithful tropicalizations of curves of degree d embedded in projective space $\mathbb{P}^r$?*

For $d = 4$ and $r = 2$ corresponding to the case of canonical curves of genus three [5] answer this question. See [7] for progress on realizability questions.

Chan and Jiradilok [6] solve Problem 1.7 and answer Question 1 when $G = K_4$. As an application of Theorem 1.1, we generalize this to all 3-regular three-connected planar graphs and when the ground field is the completion of the field of Puiseux series. More precisely, we show the following.

Let $\mathbb{K}$ be the completion of the field of Puiseux series with respect its non-trivial valuation. Let C be a smooth, proper algebraic curve over $\mathbb{K}$ and G be a 3-regular three connected planar graph whose genus is equal to that of C. Suppose that G is the combinatorial graph underlying some skeleton of the Berkovich analytification C^{an} of C.

Theorem 1.8. *There exists a canonical embedding of C that specializes to the schön embedding of X_G and this canonical embedding admits a weakly faithful tropicalization.*

The proof of Theorem 1.8 and Corollary 1.9 involves deforming the tropical curve into the tropicalization of a canonical curve such that the topology is preserved. As a corollary to Theorem 1.8, see Section 4 for more details.

Corollary 1.9. *Given a 3-regular, three-connected planar graph G, there exists a canonical curve whose Berkovich skeleton is G that admits a weakly faithful tropicalization.*

Acknowledgements: We thank to Bo Lin, Ralph Morrison and Bernd Sturmfels very much, this work has benefited significantly from the several discussions we had with them. We also thank Omid Amini, Erwan Brugalle and Lorenzo Fantini for helpful discussions.

2 A Brief Interlude into Tropical Projective Space

Tropical graph curves live in *tropical projective space*. Hence, we start by briefly recalling tropical projective space, we refer to [10, Chapter 6, Section 2] for a detailed discussion. Analogous to its classical counterpart, tropical projective space in n-dimensions $\mathbb{TP}^n$ can be constructed in different ways, we describe the one via compactification here.

Consider the hyperplane $H_0 = (1, \ldots, 1)^{\perp}$ where $(1, \ldots, 1) \in \mathbb{R}^{n+1}$ as a model for the tropical torus in n-dimensions. Each point in H_0 is a representative of an orbit of tropical multiplication of $\mathbb{R}$ on $\mathbb{R}^{n+1}$ via $\lambda \odot (q_0, \ldots, q_n) = (\lambda + q_0, \ldots, \lambda + q_n)$.

Compactify it with the $(n+1)$-coordinate hyperplanes. It is convenient to think of these hyperplanes $F_1, \ldots, F_{n+1}$, say as living at "infinity". In particular F_i is the intersection of the affine copy of the hyperplane $(0, \ldots, 0, \underbrace{1}_{i}, 0, \ldots, 0)^{\perp} \cap H_0$ at "infinity". Hence, $\mathbb{TP}^n$ is homeomorphic to the n-simplex. This identification is particularly useful for visualization purposes.

2.1 Tropicalizing into $\mathbb{TP}^n$

We recall tropicalization of a subvariety of projective space into tropical projective space. Based on the context, we tropicalize with respect to the trivial or a non-trivial valuation (in Section 3 with respect to a trivial valuation and in Section 4 with respect to a non-trivial one).

Given a homogenous polynomial $f \in \kappa[x_1, \ldots, x_{n+1}]$, suppose that $f = m \cdot g$ where m is a monomial and $g \in \kappa[x_1, \ldots, x_{n+1}]$ has no monomial factors. The tropicalization $\text{tropproj}(f)$ of f into $\mathbb{TP}^n$ is defined as the union of the tropicalization of g into H_0 (with respect to a valuation on κ) and the facets F_j for each variable x_j that divides m.

For example, if $f = (x_1^2 + x_1x_2 + x_1x_3)$ then $m = x_1$ and $g = (x_1 + x_2 + x_3)$. The tropicalization of f into $\mathbb{TP}^2$ is the union of the tropicalization of g into H_0 and the facet of $\mathbb{TP}^2$ corresponding to x_1. In contrast, the tropicalization of f into H_0 is just the tropicalization of g into H_0.

The tropicalization $\text{tropproj}(I)$ of a graded ideal I into $\mathbb{TP}^n$ is defined as:

$$\text{tropproj}(I) = \cap_{f \in I} \text{tropproj}(f)$$

A subset B of I is called a *tropical basis* for I if $\cap_{f \in B} \text{tropproj}(f) = \text{tropproj}(I)$. Every graded ideal I has a finite tropical basis [10, Theorem 2.6.5].

If the valuation underlying the tropicalization is non-trivial then the Fundamental theorem of tropical geometry holds i.e., the closure of image of the projective variety into $\mathbb{TP}^n$ under the valuation map coincides with the tropicalization of its defining ideal [10, Theorem 6.2.15].

2.2 Linear Subspaces of Tropical Projective Space:

We primarily encounter linear subspaces of $\mathbb{TP}^n$, in particular lines.

Definition 2.1. *A k-dimensional tropical linear subspace in $\mathbb{TP}^n$ is defined as the tropicalization* $\text{tropproj}(I)$ *of the defining ideal I of a k-dimensional subspace of $\mathbb{P}^n$.*

In order to tropicalize I, apriori knowing a tropical basis of I is particularly useful. Tropical bases for I have been studied in detail and a linear space has a tropical basis of linear forms [10, Proposition 4.1.6]. The linear subspaces we encounter in this paper are all defined by a set of linear forms with mutually disjoint support. For instance, the ideal $L_{v_2} = \langle x_{F_1} + x_{F_4} + x_{F_3}, x_{F_2}, x_{F_5} \rangle$ that appeared in the canonical embedding of $X_{\mathcal{C}}$ in the introduction. Such a linear subspace has a particularly simple tropical basis.

Lemma 2.2. *Suppose that I is generated by linear forms $\ell_1, \ldots, \ell_r$ such that their supports are mutually disjoint. Then the set $\{\ell_1, \ldots, \ell_r\}$ is a tropical basis for I.*

Proof. Let I_1 be the vector space of linear forms of I. Since I has a tropical basis of linear forms, if suffices to show that $\cap_{i=1}^r \text{tropproj}(\ell_i) \subseteq \cap_{\ell \in I_1} \text{tropproj}(\ell)$. Any linear form $\ell \in I_1$ is a κ-linear combination of $\ell_1, \ldots, \ell_r$. Let S be the subset of $\{\ell_1, \ldots, \ell_r\}$ with non-zero coefficients in this linear combination and let S_m be the set of monomials in S.

Because the supports of $\ell_1, \ldots, \ell_r$ are mutually disjoint, $\cap_{i \in S} \text{tropproj}(\ell_i) \subseteq \text{tropproj}(\ell) \cap \cap_{m \in S_m} \text{tropproj}(m)$. Hence, $\cap_{i=1}^{r} \text{tropproj}(\ell_i) \subseteq \cap_{\ell \in I_1} \text{tropproj}(\ell)$. □

In order to draw a sub-variety of tropical projective space with the unit simplex as a model, for each face f of dimension d, consider a system of homeomorphisms (not linear in general) from $\mathbb{R}^d$ to the relative interior of f that are compatible with each other [10, Section 6.2] and use this to transfer co-ordinates from Euclidean space to the unit simplex. Indeed, this induces a topology on $\mathbb{TP}^n$ and any subvariety of $\mathbb{TP}^n$. This is the topology on $\mathbb{T}_G$ underlying the definition of tropical graph curves. In this paper, we are primarily interested in the topology of subvarieties (in particular, arrangements of lines) of $\mathbb{TP}^n$ and the figures are all only accurate up to topology.

3 Existence and Realizability of Tropical Graph Curves

In this section, we prove the Existence and Realizability theorems (Theorem 1.1 and Theorem 1.2). We begin by generalizing the schön embedding of the cube described in the introduction to arbitrary (3-regular) three-connected planar graphs.

3.1 Schön Embedding

Let G be a 3-regular, three-connected planar graph of genus g. Given a planar embedding of the graph of genus g, label the interior faces of the planar embedding by variables and let x_F be the variable corresponding to the face F. Hence, there are g variables. Let R be the graded κ-algebra generated by x_F where F ranges over all the interior faces of the planar embedding of G and identify $\mathbb{P}^{g-1}$ with $\text{Proj}(R)$. In the following, we associate to G a line arrangement in $\mathbb{P}^{g-1}$. For each vertex $v \in G$, associate an ideal L_v defined by a collections of linear forms as follows:

1. If v is an exterior vertex then, let L_v be the ideal generated by x_F over all faces F not incident on v. Note that there are precisely $g-2$ such faces.

2. If v is an interior vertex then, let L_v be the ideal generated by x_F over all faces F not incident on v (there are $g-3$ such faces) and the sum of the variables corresponding to the three faces incident on v.

In both cases above, L_v defines a line in $\mathbb{P}^{g-1}$. We refer to this line arrangement as the *schön embedding* of the graph curve X_G (this is a canonical embedding of X_G, see Corollary 3.3).

The line L_v for an exterior vertex is generated by $(g-2)$ variables and the line L_v for an interior vertex is generated by $(g-3)$ variables and one linear form given by the sum of the three remaining variables. The tropicalization of lines corresponding to exterior vertices is not branched whereas the tropicalization of lines corresponding to interior vertices contains one branch of valence three.

We refer to the former as **type I** lines and the latter as **type II** lines. For a tropical line of type II, suppose that $x + y + z$ is the associated linear form, the three branches of this line are labelled by pairs of variables (x, y), (y, z) and (x, z) according to the pair that is minimized along that branch and the point of intersection between these three branches is called the *branch point* of the line.

Bayer and Eisenbud [3, Section 6] work with schön embedding in terms of the minimal generators of its defining ideal. Showing that the schön embedding yields a weakly faithful tropicalization by directly tropicalizing the minimal generators seems difficult, its primary decomposition given by the schön embedding is more amenable from a tropical point of view. Note that the primary decomposition of the defining ideal does not depend on the ground field (as long as it is algebraically closed) and the tropicalization of the defining ideal is equal to the union of the tropicalization of its irreducible components (both over the trivial and non-trivial valuations).

Though the schön embedding is presented in terms of a planar embedding of G, it does not depend on the choice of the planar embedding. It has the following simple description in terms Stanley-Reisner varieties.

Let M be the dual complex of the three-dimensional polytope P associated to G and let F be the set of facets of P (the facets of P are in bijection with the vertices of M). The schön embedding is the Stanley-Reisner variety of M intersected with the hyperplane $\sum_{f \in F} x_f = 0$ and this hyperplane section is reduced.

Proposition 3.1. (**Schön Embedding in terms of Stanley-Reisner Variety)** A minimal generating set $\mathcal{G}$ of the defining radical ideal I_{sch} of the schön embedding is the set of minimal monomial generators of the Stanley-Reisner ideal of M with the outerface face x_e replaced by $-\sum_{f \in F, f \neq e} x_f$.

Proof. The projective schemes defined by both the Stanley-Reisner ideal of M, [3, Corollary 6.2] and the schön embedding of X_G are both reduced and are arrangements of $2g - 2$ lines in $\mathbb{P}^{g-1}$. Hence, to show that they are equal, it suffices to show that every minimal generator of the Stanley-Reisner ideal of M is contained in the ideal L_v for every vertex $v \in G$.

Recall that every element in $\mathcal{G}$ corresponds to a minimal non-face of M. An element g of $\mathcal{G}$ is either purely a product of variables or is divisible by $\sum_{f \in F, f \neq e} x_f$. Suppose that g is a product of variables. Consider L_v where v is an exterior vertex of the planar embedding of G. Suppose that g is not contained in L_v. Suppose that $F_1,\ F_2$ are the two faces incident on v then g must be one of $x_{F_1},\ \ x_{F_2}$ or $x_{F_1} \cdot x_{F_2}$. But then since F_1, F_2 share an edge in M, this is not possible. A similar argument shows that g is contained in L_v where v is an interior vertex.

On the other hand, if g is divisible by $-\sum_{f \in F, f \neq e} x_f$ then suppose that L_v is an interior vertex then $\sum_{f \in F, f \neq e} x_f \in L_v$ and hence, $g \in L_v$. Otherwise, if L_v is an exterior vertex then since $g \neq -\sum_{f \in F, f \neq e} x_f$, it must divide x_F where x_F where F is a face other than x_{F_1} and x_{F_2} where $\{F_1, F_2\}$ are the two faces incident on v. This is because $F_1,\ F_2$ and the outer face of the planar embedding of G form a face of M. Hence, $g \in L_v$.

□

Remark 3.2. *Bayer and Eisenbud [3] use this description in terms of Stanley-Resiner varieties to compute the Clifford index of the underlying graph curve.*

As an immediate consequence of Theorem 3.1 and [3, Corollary 6.2], we obtain.

Corollary 3.3. *The schön embedding is a canonical embedding of X_G.*

Theorem 3.4. (**Tropical Basis of the Schön Embedding**) The minimal generating set $\mathcal{G}$ presented in Theorem 3.1 is a tropical basis.

Proof. Let $\mathcal{G} = \{g_1, \ldots, g_r\}$. For the sake of contradiction, suppose that $\mathcal{G}$ is not a tropical basis then there is a point in $p \in \cap_{j=1}^{r} \text{tropproj}(g_j)$ that is not contained in the tropicalization of I_{sch}. Since, the generators are all products of linear forms, this implies that there is a choice of linear forms $\ell_1, \ldots, \ell_r$ such that $\ell_j | g_j$ for each j from one to r and such that $p \in \cap_{j=1}^{r} \text{tropproj}(\ell_j)$.

Consider the ideal generated by $\ell_1, \ldots, \ell_r$, This ideal contains the defining ideal of X_G. Hence, the subvariety defined by it is contained in X_G. This subvariety is either a point or an irreducible component of the graph curve. By Lemma 2.2, the set $\{\ell_1, \ldots, \ell_r\}$ is a tropical basis. Hence, p is contained in the tropicalization of $\langle \ell_1, \ldots, \ell_r \rangle$ and hence, in the tropicalization of the graph curve. This is a contradiction. □

Example 3.5. *Let $\mathcal{C}$ be the one-skeleton of the three-dimensional cube. The minimal generating set of the vanishing ideal of $X_{\mathcal{C}}$ produced by Theorem 3.1 is the following:*

$$\left\langle x_{F_2} x_{F_4}, x_{F_3} x_{F_5}, x_{F_1}^2 + x_{F_1} x_{F_2} + x_{F_1} x_{F_3} + x_{F_1} x_{F_4} + x_{F_1} x_{F_5} \right\rangle$$

Hence, $X_{\mathcal{C}}$ is cut-out by three (degenerate) quadrics in $\mathbb{P}^4$ and according to Theorem 3.4 these quadrics also form a tropical basis for $X_{\mathcal{C}}$. Furthermore, the curve $X_{\mathcal{C}}$ is a complete intersection. But in general, canonical embeddings of graph curves are not complete intersections.

Finally, we note that the dual of a canonical embedding of X_G is not necessarily always G. A 3-regular graph G is called *earnest* if the dual graph of a canonical embedding of X_G is equal to G.

Example 3.6. *Consider the graph G as shown Figure 3. As shown in [13, Remark 1.2], no arrangement of lines in projective space has G as its dual graph. Hence, G is not earnest.*

As an immediate consequence of Theorem 3.1 and [3, Corollary 6.2] we obtain.

Corollary 3.7. *Let G be a simple, 3-regular three-connected planar graph, the schön embedding has dual graph G. Hence, G is an earnest graph.*

More generally, Bayer and Eisenbud [3, Proposition 2.5, Item ii] show that a (3-regular) graph is earnest if and only it is three-connected.



Figure 3: Non-Earnest Graph



Figure 4: ΔY Transformation

3.2 Tropicalization of the Schön Embedding

The proofs of the Existence and Realizibility theorems adapt a proof idea of Steinitz theorem. Hence, we begin by recalling Steinitz theorem ([15, Chapter 4]) and one of its proofs.

Theorem 3.8. (**Steinitz Theorem**) *A planar graph is the one-skeleton of the graph of a three-dimensional polytope if and only if it is three vertex connected.*

A key ingredient in the proof is a ΔY transformation i.e., replace a Δ-subgraph (a triangular face) by a Y-subgraph, as shown in Figure 4. A simple ΔY transformation is a ΔY transformation followed by replacing multi edges created by this transformation by simple edges and contracting edges to eliminate valence two vertices. The next step is to show that every three-connected planar graph can be obtained from K_4 by a sequence of simple ΔY transformations. The graph K_4 is realizable via a tetrahedron and the fact that if a planar graph is realizable by a three-dimensional polytope then the resulting graph after a ΔY transformation is also realizable concludes the proof.

We adapt this framework to tropicalizations of schön embeddings instead of one-skeleta of three-dimensional polytopes. Since, we are concerned with 3-regular three-connected planar graphs rather than arbitrary three-connected planar graphs, we employ a sequence of "local operations" that transform a 3-regular three-connected planar graph to K_4 while keeping the property 3-regular invariant. We perform the following two steps:

1. A ΔY transformation.

2. A *contraction-elongation* operation shown in Figure 5. We say that the operation is performed on the edge e_1.



Figure 5: Contraction-Elongation Operation



Figure 6: Two Exterior Vertices Sharing an Interior Face

We also employ the following characterization of trivalent, three-connected planar graphs. Recall that a bridge is an edge whose deletion disconnects the graph.

Lemma 3.9. *Let G be a simple, 3-regular, bridgeless, planar graph. The graph G is three edge connected if and only if any planar embedding of G is such that no two non-adjacent exterior vertices share an interior face.*

Proof. ($\Rightarrow$) Suppose that there is a planar graph of G such that there exist two non-adjacent exterior vertices u and v, say that share an interior face. Consider the edges e_1 and e_2 incident on u and v, respectively that are shared by this interior face and the exterior face. Deleting this two edges e_1 and e_2 will disconnect the graph. Hence, the graph is not three edge connected.

($\Leftarrow$) Conversely, suppose that G is not a three edge connected graph. Since, G is bridgeless, it is two edge connected. Suppose that deletion of edges e_1 and e_2 disconnects G. Consider a planar embedding of G. The edges e_1 and e_2 cannot share a vertex v since this implies that the other edge incident on v is a bridge. Hence, the edges e_1 and e_2 bind an interior face F of G, see Figure 6. The vertices v_1 and u_2 are not adjacent and share an interior face.

□

Remark 3.10. *Since three edge connectivity is equivalent to three vertex connectivity for 3-regular graphs [3, Lemma 2.6], we can replace edge connectivity with vertex connectivity in Lemma 3.9.*

Lemma 3.11. *Every 3-regular, three-connected planar graph G can be transformed to K_4 by a sequence of ΔY and contraction-elongation operations such that the graph at each step remains a 3-regular, three-connected planar graph.*

Proof. Suppose that G has genus three, then it is a K_4 and there is nothing to prove. Otherwise, the genus of G is at least four. Consider a planar embedding of G, we perform the following operations:

1. Suppose that G has a Δ then perform a Δ to Y operation on it.
2. Otherwise, since the graph is trivalent, it has an interior face. Take an interior face F of the minimum length, k say. It has at least $k-3$ interior edges (edges not incident on the exterior face). Perform a sequence of contraction-elongation operations on any $k-3$ interior edges. This results in (at least) one Δ.
3. Perform a Δ to Y-transformation on one of these Δs.

We first show that every graph produced by this procedure is simple, 3-regular and three-connected. Any contraction-elongation operation on an interior edge does not create a bridge, suppose it does then this implies that the bridge is the edge e_2, see Figure 5. But this implies that the original graph can be disconnected by deleting two edges. Furthermore, it does not alter the exterior face and the set of faces incident on each exterior vertex remains unaltered. Hence, no two non-adjacent exterior vertices can share an interior face after the operation. By Lemma 3.9, the resulting graph remains three-connected.

We now show that the graph resulting from a ΔY transformation remains simple, 3-regular and three-connected. For a multiple edge to occur during a ΔY transformation, the subgraph induced by the vertices of the Δ must be a K_4 minus an edge as shown in Figure 7. But this contradicts the three-(edge) connectivity of the graph on which the operation is performed since deleting the edges e_1 and e_2 would disconnect the graph. By the proof of Steinitz theorem, the graph remains three-connected. Hence, the result G' is a simple, 3-regular graph and its genus $g(G') = g(G) - 1$.

Repeat Operation 2 till the genus of the resulting graph is three. This graph must be K_4 since it is the only 3-regular, simple graph of genus three.

□

Next, we show that the tropicalization of the schön embedding of X_{K_4} is weakly faithful (Lemma 3.13). We show that Y to Δ operations and contraction-elongation operations preserve this property (Lemmas 3.14 and 3.15). We conclude that the schön embedding of every three-connected planar graph admits a weakly faithful tropicalization.



Figure 7: Forbidden Subgraph

In the following, we employ a property that relates the homeomorphism that realizes a weakly faithful tropicalization to the graph G. Before stating this property, we note that for a 3-regular planar graph, every exterior vertex has precisely one interior vertex adjacent to it.

Definition 3.12. *Let G be a 3-regular connected graph. A weakly-faithful tropicalization of X_G is said to be* well structured *(with respect to a planar embedding of G) if there exists a homeomorphism from the modification of G to the tropicalization of X_G that satisfies the following properties:*

- *For an interior vertex v, its image in the tropicalization is the branch point of L_v.*
- *For an exterior vertex v, its image in the tropicalization is the intersection between L_v and L_u where u is the unique interior vertex adjacent to it.*

Lemma 3.13. *The schön embedding of K_4 admits a weakly faithful tropicalization and this tropicalization is well structured.*

Proof. Consider the embedding of K_4 with a triangle exterior face and one vertex in the interior. The schön embedding of K_4 is as follows: $L_{v_1} = x_{F_1}$, $L_{v_2} = x_{F_2}$, $L_{v_3} = x_{F_3}$, $L_{v_4} = (x_{F_1} + x_{F_2} + x_{F_3})$ where F_1, F_2 and F_3 are the three interior faces. Its tropicalization is as shown in Figure 8. This is weakly faithful and well structured. □

Lemma 3.14. *Let G be a simple, 3-regular, three-connected planar graph and let G' be a graph obtained after a Y to Δ-transformation. If the tropicalization of the schön embedding of X_G is weakly faithful and well structured, then the tropicalization of the schön embedding of $X_{G'}$ is also weakly faithful and well structured.*

Proof. Suppose that the Y has an exterior vertex v as its trivalent vertex as shown in Figure 9. Let F_1 and F_2 be the two faces incident on v and let F_n be the new face created by the



Figure 8: Tropicalization of the Schön Embedding of K_4

.



Figure 9: Y to Δ on an Exterior Vertex and its Tropicalization



Figure 10: Y to Δ on an Interior Vertex and its Tropicalization

transformation. We study the change in the tropicalization of the schön embedding under this transformation.

The line L_v is given by $\langle x_F |\ F \notin \{F_1, F_2\}\rangle$ is replaced by three lines L_{v_1}, L_{v_2} and L_{v_3}. They are given by the

$$\begin{aligned} L_{v_1} &= \langle x_F |\ F \notin \{F_1, F_n\}\rangle, \\ L_{v_2} &= \langle x_F |\ F \notin \{F_2, F_n\}\rangle \\ L_{v_3} &= \langle x_{F_n} + x_{F_1} + x_{F_2},\ x_F |\ F \notin \{F_n, F_1, F_2\}\rangle \end{aligned}$$

The changes in the tropicalization is shown in the Figure 9.

Suppose that the graph Y is centered at an interior vertex v as shown in Figure 10 and let F_1, F_2 and F_3 be the three faces incident on it. Suppose that F_n is the face created by the Δ and v_1, v_2 and v_3 be the newly created vertices. The line L_v is given by $\langle x_{F_1} + x_{F_2} + x_{F_3}, x_F |\ F \notin \{F_1, F_2, F_3\}\rangle$ and the lines L_{v_1}, L_{v_2} and L_{v_3} are given by:

$$\begin{aligned} L_{v_1} &= \langle x_{F_1} + x_{F_2} + x_{F_n}, x_F |\ F \notin \{F_1, F_2, F_n\}\rangle, \\ L_{v_2} &= \langle x_{F_2} + x_{F_3} + x_{F_n} x_F |\ F \notin \{F_2, F_3, F_n\}\rangle \\ L_{v_3} &= \langle x_{F_1} + x_{F_3} + x_{F_n},\ x_F |\ F \notin \{F_1, F_3, F_n\}\rangle \end{aligned}$$

The changes in the tropicalization is shown in the Figure 10.

In the following, we show that the tropicalization of the schön embedding is homeomorphic to G' (up to a modification) and is well structured.

Consider a homeomorphism ϕ realizing the well structured weakly faithful tropicalization from a modification G_{mod} of G to the tropicalization of schön embedding. Let S be the subgraph induced by v, the trivalent vertex of Y (on which the ΔY operation is performed). Note that S is the graph with one vertex and no edges. Similarly, let S' be the subgraph of G' respectively induced by the resulting Δ.



Figure 11: Tropicalization of the Contraction-Elongation Operation

Let $\mathcal{A}_{S'}^{\text{trop}}$ be the tropicalization of the (sub)-arrangement of the schön arrangement of $X_{G'}$ formed by lines L_v where $v \in S'$. There is a homeomorphism ψ from a modification of S' to $\mathcal{A}_{S'}^{\text{trop}}$ that is well structured.

Let $\phi|(G_{\text{mod}} \setminus S)$ be the restriction of ϕ to $G_{\text{mod}} \setminus S$. Let G'_{mod} be the modification of G' induced by (given by the disjoint union of) the modification of S' corresponding to ψ and the modification of G' induced by ϕ. More precisely, there is a canonical isomorphism from $G \setminus S$ to $G' \setminus S'$ and the modification G_{mod} induces a modification G'_{mod} via this isomorphism. Consider a homeomorphism ϕ' from G'_{mod} to the tropicalization of the schön embedding of $X_{G'}$ that agrees with $\phi|(G_{\text{mod}} \setminus S)$ (via the canonical isomorphism between $G' \setminus S'$ and $G \setminus S$) when restricted to $G' \setminus S'$ and with ψ when restricted to S'. This is a homeomorphism from a modification of G' to the tropicalization of the schön embedding of $X_{G'}$. Furthermore, it is well structured since both ϕ and ψ respect the property of well structuredness.

□

Lemma 3.15. *Let G be a simple, 3-regular, three-connected, planar graph and let G' be a graph obtained after a contraction-elongation operation along an interior edge e_1 (an edge not incident on the outer face). If the tropicalization of the schön embedding of X_G is weakly faithful and well structured, then the tropicalization of the schön embedding of $X_{G'}$ is also weakly faithful and well structured.*

Proof. Suppose that v_1 and v_2 are vertices incident on e_1 and suppose that w_1 and w_2 vertices incident on the edge e_2 that is created by the operation, see Figure 5. The lines L_{v_1} and L_{v_2} are given by equations $\langle x_{F_1} + x_{F_3} + x_{F_4}, x_F |\ F \notin \{F_1, F_3, F_4\}\rangle$ and $\langle x_{F_2} + x_{F_3} + x_{F_4}, , x_F |\ F \notin \{F_2, F_3, F_4\}\rangle$. Similarly, the lines L_{w_1} and L_{w_2} are given by equations $\langle x_{F_1} + x_{F_2} + x_{F_3}, x_F |\ F \notin \{F_1, F_2, F_3\}\rangle$ and $\langle x_{F_1} + x_{F_2} + x_{F_4}, x_F |\ F \notin \{F_1, F_2, F_4\}\rangle$. The tropicalization changes as shown in the Figure 11. In the following, we note that the tropicalization is homeomorphic to G' up to modification and is well structured.

The argument is similar to the one in the proof of Lemma 3.14. Let S and S' be the subgraphs of G and G' induced by $\{v_1, v_2, u_1, u_2, u_3, u_4\}$ and $\{w_1, w_2, u_1, u_2, u_3, u_4\}$ respectively. Note that there is a canonical isomorphism between $G \setminus S$ and $G' \setminus S'$.

Let ϕ be a homeomorphism from a modification G_{mod} of G that is weakly faithful and well structured. Let ψ be a homeomorphism from a modification S'_{mod} to the arrangement formed by the tropicalization of the lines L_v where $v \in \{w_1, w_2, u_1, u_2, u_3, u_4\}$ takes the vertices w_1 and w_2 to the corresponding branch points of $\text{trop}(L_{w_1})$ and $\text{trop}(L_{w_2})$ (see Figure 11).

There is a homeomorphism ϕ' from a modification G'_{mod} of G' (induced by ψ and by ϕ via the canonical isomorphism from G to G') to the tropicalization of the schön embedding of $X_{G'}$. This agrees with ψ when restricted to S' and with $\phi|(G_{\text{mod}} \setminus S)$ when restricted to $G'_{\text{mod}} \setminus S'$ via the canonical isomorphism from $G \setminus S$ and $G' \setminus S'$. This homeomorphism is well structured since both ϕ and ψ respect the property of well structuredness.

□

Theorem 3.16. *Let G be a three-connected planar graph. The schön embedding of G admits a weakly faithful tropicalization into tropical projective space and this tropicalization is well structured.*

Proof. By Lemma 3.11, there exists a sequence of Y to Δ and contraction-elongation operations to transform K_4 to G. By Lemma 3.13, the tropicalization of the schön embedding of K_4 is weakly faithful and well structured. By Lemmas 3.14 and 3.15, we conclude that the tropicalization of the schön embedding of G is weakly faithful and is well structured. □

Example 3.17. *Consider the one-skeleton of the three dimensional cube. A sequence of transformations following Lemma 3.11 to transform it into K_4 is shown in Figure 12. A sequence is the one-skeleton of the following polytopes:*

cube → triangular prism sliced at a vertex → triangular prism → tetrahedron.

The tropical counterpart of the sequence is shown in Figure 13.

4 Lifting Graph Curves

In this section, we deform the schön embedding to construct weakly faithful tropicalizations of canonical curves. Let G be a 3-regular, three-connected planar graph of genus g. Suppose that C is a smooth curve of genus g whose Berkovich analytification has a skeleton whose underlying graph is G, we first show that the schön embedding of X_G deforms to a canonical embedding of C.

Let Γ be the value group of $\mathbb{K}$, R be the ring of integers of $\mathbb{K}$ and κ be the residue field of $\mathbb{K}$.

Lemma 4.1. *Let C be a smooth, proper algebraic curve over $\mathbb{K}$ and G be a 3-regular, three connected planar graph whose genus is equal to that of C. Suppose that G is the graph underlying a skeleton of the Berkovich analytification C^{an} of C, then any canonical embedding ω_G of the graph curve X_G deforms to a canonical embedding of C.*



Figure 12: Example: Sequence of Polytopes



Figure 13: Example: Corresponding Tropical Graph Curves

Proof. From [4, Chapter 4], we know that C has a semistable model $\mathfrak{X}$ (defined over $\mathrm{Spec}(R)$) such that the special fiber of $\mathfrak{X}$ is the graph curve X_G (over κ). The arithmetic surface $\mathfrak{X}$ has a canonical bundle $\Omega_{\mathfrak{X}}$ that over the generic fiber restricts to the canonical bundle Ω_X of X and over the special fiber restricts to the canonical bundle Ω_{X_G} of X_G [9].

Choose a basis $\mathfrak{B}$ for $H^0(\Omega_{\mathfrak{X}})$ and let $\mathfrak{B}_0$ be the reduction of $\mathfrak{B}$ to a basis for $H^0(\Omega_{X_G})$. Given a canonical embedding of X_G, defined by a basis $\mathfrak{B}'_0$ of $H^0(\Omega_{X_G})$. Let M_0 be the element in $\mathrm{GL}(g,\kappa)$ that takes $\mathfrak{B}_0$ to $\mathfrak{B}'_0$. Consider any lifting of M_0 to an element M of $\mathrm{GL}(g,R)$ (such that an element always exists since the reduction map induces a surjection from $\mathrm{GL}(g,R)$ to $\mathrm{GL}(g,\kappa)$). Consider the basis for $H^0(\Omega_{\mathfrak{X}})$ obtained by applying M to $\mathfrak{B}$. When restricted to the generic fiber this yield a canonical embedding of C that has the required property. □

Corollary 4.2. *Let C be a smooth, proper algebraic curve over $\mathbb{K}$ and let G be a 3-regular, three connected graph whose genus is equal to that of C. Suppose that G is a skeleton of C^{an} then the schön embedding of X_G deforms to a canonical embedding of C.*

The canonical ideal of C constructed in Corollary 4.2 is defined over R, the ring of integers of $\mathbb{K}$. We call such a (any) deformation of the schön embedding of X_G a *schön embedding* of C. Note that C can have multiple schön embeddings. We now show that any schön embedding admits a weakly faithful tropicalization.

Theorem 4.3. *Let $\mathbb{K}$ be the completion of the field of Puiseux series with respect to its non-trivial valuation. Let C be a smooth, proper algebraic curve over $\mathbb{K}$ and G be a 3-regular three connected planar graph whose genus is equal to that of C. Suppose that G is the combinatorial graph underlying a skeleton of the Berkovich analytification C^{an} of C, then a schön embedding of C admits a weakly faithful tropicalization.*

Proof. Consider a schön embedding of C. Let $\mathbb{K}[x_1,\ldots,x_g]$ be the co-ordinate ring of $\mathbb{P}^{g-1}$ and let I_C be the defining ideal of this schön embedding of C. The initial form with respect to a weight vector $\mathbf{w}=(w_1,\ldots,w_{g+1})$ is defined as:

$$\mathrm{in}_{\mathbf{w}}(f) = \sum_{\alpha} c_\alpha \mathbf{x}^\alpha, \text{ over all terms } \alpha \text{ is the support of } f \text{ such that } w_{g+1}\mathrm{val}(c_\alpha) + (w_1,\ldots,w_g)\cdot\alpha \text{ is minimized.}$$

Homogenize I_C with respect to a weight vector with weight zero at each variable and weight minus one for the uniformizing parameter t. Let h be the homogenizing variable. This results in a flat family over h. The fiber over $h=1$, is the ideal I_C and the fiber over $h=0$ is the schön embedding of X_G.

Consider the tropicalization of this family with respect to the non-trivial valuation val on $\mathbb{K}$ into $\mathbb{TP}^{g-1}\times\mathbb{R}\cup\{\infty\}$. By the theorem Bieri-Groves, the tropicalization of this family is a polyhedral surface.

Let us briefly describe the structure of this polyhedral surface. Let y be the variable corresponding to h. The fiber over $y=k$ corresponds to tropicalizing the surface at $h=t^k$. By construction, two fibers over non-negative k_1 and k_2 in $\mathbb{R}$ are a scaling of each other. The fiber over $y=\infty$ gives the tropicalization of the schön embedding over $\mathbb{K}$ and is homeomorphic to G. Hence, the fibers over y for $y\neq -1$ are homeomorphic to a

modification of G. By Theorem 3.16, the tropicalization of the schön embedding of X_G has genus g. Furthermore, the genus of each fiber over $y \geq 0$ of the family is g.

Hence, $\mathrm{trop}(C)$ has genus g. By [2, Section 2.3], there exists a continuous map from the Berkovich analytification of C to $\mathrm{trop}(C)$. Hence, the genus of Berkovich analytification is equal to g and this continuous map is a homeomorphism. Hence, the tropicalization of C is weakly faithful. □

Remark 4.4. *Note that in the proof of Theorem 4.3, it is more natural to tropicalize the schön embedding X_G over the non-trivial valuation over $\mathbb{K}$ rather than the trivial valuation since X_G is realized as a limit of smooth curves over $\mathbb{K}$. However, the tropicalizations with respect to* val *and the trivial valuation are both equal.*

For every 3-regular, three connected graph G of genus g, there exists a smooth curve of genus g with G as a skeleton [1, Appendix]. Hence, we answer Question 1 for 3-regular three-connected planar graphs.

Corollary 4.5. *Let $\mathbb{K}$ be the completion of the field of Puiseux series with respect its non-trivial valuation. Let G be a 3-regular three-connected planar graph of genus g. There exists a canonical curve C with Berkovich skeleton whose underlying graph is G and whose tropicalization is weakly faithful.*

Example 4.6. *Let $\mathcal{C}$ be the one-skeleton of the three-dimensional cube. Consider the graph curve $X_{\mathcal{C}}$ over $\mathbb{K}$ (as in Theorem 4.3) in its schön embedding. It is defined by the ideal*

$$\langle x_{F_2}x_{F_4}, x_{F_3}x_{F_5}, x_{F_1}^2 + x_{F_1}x_{F_2} + x_{F_1}x_{F_3} + x_{F_1}x_{F_4} + x_{F_1}x_{F_5}\rangle \tag{1}$$

Consider three smooth quadratics in $\mathbb{K}[x_{F_1}, x_{F_2}, x_{F_3}, x_{F_4}, x_{F_5}]$ defined the ring of integers of $\mathbb{K}$ that deforms the minimal generators.

For instance, consider the three quadrics such that homogenizing with respect to the weight vector $(0,0,0,0,0,-1)$ where the first five coordinates correspond to the variables $x_{F_1}, \ldots, x_{F_5}$ and the last coordinate corresponds to t leads to the following:

$$q_1 = t^{100}x_{F_1}^2h^{100} + t^{50}x_{F_2}^2h^{50} + t^{75}x_{F_3}h^{75} + t^{17}x_{F_4}^2h^{17} + t^{15}x_{F_5}^2h^{15} + t^{16}x_{F_1}x_{F_2}h^{16} + t^{27}x_{F_1}x_{F_3}h^{27} + t^{18}x_{F_1}x_{F_4}h^{18} + t^{78}x_{F_1}x_{F_5}h^{78} + t^{-34}x_{F_2}x_{F_3}h^{34} + t^{150}x_{F_2}x_{F_5}h^{150} + t^{52}x_{F_3}x_{F_4}h^{52} + t^{189}x_{F_3}x_{F_5}h^{189} + t^{19}x_{F_4}x_{F_5}h^{19} + x_{F_2}x_{F_4}\ ,$$

$$q_2 = t^{156}x_{F_1}^2h^{156} + t^{151}x_{F_2}^2h^{151} + t^{176}x_{F_3}^2h^{176} + t^{117}x_{F_4}^2h^{117} + t^{5}x_{F_5}^2h^{5} + t^{61}x_{F_1}x_{F_2}h^{61} + t^{7}x_{F_1}x_{F_3}h^{7} + t^{81}x_{F_1}x_{F_4}h^{81} + t^{82}x_{F_1}x_{F_5}h^{82} + t^{43}x_{F_2}x_{F_3}h^{43} + t^{67}x_{F_2}x_{F_4}h^{67} + t^{57}x_{F_2}x_{F_5}h^{57} + t^{87}x_{F_3}x_{F_4}h^{87} + t^{92}x_{F_4}x_{F_5}h^{92} + x_{F_3}x_{F_5},$$

$$q_3 = t^{111}x_{F_2}^2h^{111} + t^{166}x_{F_3}^2h^{166} + t^{171}x_{F_4}^2h^{171} + t^{5}x_{F_5}^2h^{5} + t^{34}x_{F_2}x_{F_3}h^{34} + t^{75}x_{F_2}x_{F_4}h^{75} + t^{41}x_{F_2}x_{F_5}h^{41} + t^{78}x_{F_3}x_{F_4}h^{78} + t^{16}x_{F_3}x_{F_5}h^{16} + t^{29}x_{F_4}x_{F_5}h^{29} + x_{F_1}^2 + x_{F_1}x_{F_2} + x_{F_1}x_{F_3} + x_{F_1}x_{F_4} + x_{F_1}x_{F_5}.$$

A key property, as required by Theorem 4.3, is that the ideal generated by q_1, q_2, and q_3 in $R[x_1, \ldots, x_5]$ reduces to schön embedding of $X_{\mathcal{C}}$ (see Equation (1)) over the residue field. These polynomials cut out a surface in $\mathbb{P}^4 \times \mathbb{A}^1$. Tropicalize it with respect to the non-trivial valuation. This results in a surface in $\mathbb{TP}^4 \times (\mathbb{R} \cup \{\infty\})$. Let w be the variable corresponding to h. The fiber over $w = r$ corresponds to the tropicalization of the curve obtained by specializing h to an element of valuation r and the fiber at infinity is $\mathrm{trop}(X_{\mathcal{C}})$.

References

[1] Matthew Baker, *Specialization of linear systems from curves to graphs*, Algebra and Number Theory **2(6)**, 613–653, 2008.

[2] Matthew Baker, Sam Payne and Joseph Rabinoff, *Nonarchimedean geometry, tropicalization, and metrics on curves*, arXiv:1104.0320, to appear in Algebraic Geometry, 2011.

[3] Dave Bayer and David Eisenbud, *Graph Curves*, Advances in Mathematics **86**, 1–40, 1991.

[4] Vladimir G. Berkovich, *Spectral Theory and Analytic Geometry over Non-Archimedean Fields*, Mathematical Surveys and Monographs of the American Mathematical Society **33**, 1990.

[5] Sarah Brodsky, Michael Joswig, Ralph Morrison and Bernd Sturmfels, *Moduli of tropical plane curves,* Research in the Mathematical Sciences **2:4**, 2015.

[6] Melody Chan and Pakawut Jiradilok, *Theta characteristics of tropical K_4-curves*, arXiv:1503.05776, 2015.

[7] Man-Wai Cheung, Lorenzo Fantini, Jennifer Park and Martin Ulirsch, *Faithful Realizability of Tropical Curves*, International Mathematics Research Notices, available at doi: 10.1093/imrn/rnv269, 2015.

[8] Andreas Gathmann and Michael Kerber, *A Riemann-Roch theorem in tropical geometry*, Mathematische Zeitschrift **259 (1)**, 217–230, 2008.

[9] Eric Katz, Joseph Rabinoff and David Zureick-Brown, *Uniform bounds for the number of rational points on curves of small Mordell–Weil rank*, arXiv:1504.00694, to appear in Duke Mathematical Journal, 2015.

[10] Diane Maclagan and Bernd Sturmfels: *Introduction to Tropical Geometry*, American Mathematical Society, Graduate Studies in Mathematics **161**, 2015.

[11] Grigory Mikhalkin and Ilia Zharkov, *Tropical curves, their Jacobians and Theta functions*, Proceedings of the International Conference on Curves and Abelian Varieties, 2007.

[12] Tim Römer and Kirsten Schmitz, *Generic tropical varieties*, Journal of Pure and Applied Algebra **216**, 140–148.

[13] Zvi Rosen, *Graph Curves*, available at https://math.berkeley.edu/ zhrosen/graphcurves.pdf, 2011.

[14] Amaury Thuillier, *Thorie du potentiel sur les courbes en gomtrie analytique non archimdienne. Applications la thorie d'Arakelov*, PhD thesis, Université Rennes, 2005.

[15] Günter Ziegler, *Lecture Notes on Polytopes*, Graduate Texts in Mathematics, Springer 1995.

Authors' address:

School of Mathematical Sciences,
Queen Mary University of London,
London, E1 4NS.
Email id: m.manjunath@qmul.ac.uk